

ANTHRAX

ALL SUSPECT CASES OF ANTHRAX MUST BE REPORTED IMMEDIATELY TO THE HUMBOLDT COUNTY PUBLIC HEALTH BRANCH:

During business hours: 707-445-6200
After hours: Sheriff's Dispatch, (ask for the Health Officer) 707-445-7251

Epidemiology:

- Anthrax can be transmitted by inhalation, ingestion, or inoculation (inhalation is the most likely during a bioterrorist attack)
- The spore form of anthrax is highly resistant to physical and chemical agents; spores can persist in the environment for years
- **Anthrax is not transmitted from person to person**

Clinical:

- Incubation period is 1-5 days (range up to 43 days)
- Inhalation anthrax presents as acute hemorrhagic mediastinitis
- Biphasic illness, with initial phase characterized by nonspecific flu-like illness followed by acute phase characterized by acute respiratory distress and toxemia (sepsis)
- Chest x-ray findings: **Mediastinal widening in a previously healthy patient in the absence of trauma is pathognomonic for anthrax**
- Mortality rate for inhalation anthrax approaches 90%, even with treatment. Shock and death within 24 – 36 hours

Laboratory Diagnosis:

- Laboratory specimens should be handled in a Biosafety Level 2 facility (e.g. California state Microbial Diseases Laboratory)
- Gram stain shows gram positive bacilli, occurring singly or in short chains, often with squared off ends (safety pin appearance). In advanced disease, a gram stain of unspun blood may be positive
- Distinguishing characteristics on culture include: non-hemolytic, non-motile, capsulated bacteria that are susceptible to gamma phage lysis
- ELISA and PCR tests are available at national reference laboratories

Patient Isolation:

- Standard barrier isolation precautions. Patients do not require isolation rooms
- **Anthrax is not transmitted person to person**

Treatment:

- Prompt initiation of antibiotic therapy is essential
- Antibiotic susceptibility testing is KEY to guiding treatment
- Ciprofloxacin (400 mg IV q 12 hr) is the antibiotic of choice for penicillin-resistant anthrax or for empiric therapy while awaiting susceptibility results
- All patients should be treated with anthrax vaccine if available; antibiotic treatment should be continued until 3 doses of vaccine have been administered (day 0, 14 and 28). If vaccine is unavailable, antibiotic treatment should be continued for 60 days.

Prophylaxis:

- If vaccine is available, all exposed persons (as determined by local and state health depts) should be vaccinated with 3 doses of anthrax vaccine (days 0, 14 and 28)

- Start antibiotic prophylaxis immediately after exposure with ciprofloxacin (500 mg po q 12 hrs) or doxycycline (100 mg po q 12 hrs). (If strain is penicillin-susceptible, therapy can be modified to penicillin or amoxicillin.)
- Antibiotic prophylaxis should be continued until 3 doses of vaccine have been administered; if vaccine is unavailable, antibiotics should be continued for 60 days.